

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 June 2004 (24.06.2004)

PCT

(10) International Publication Number
WO 2004/053158 A1

(51) International Patent Classification⁷:
C07K 14/72

C12Q 1/68 //

(74) Agent: OY JALO ANT-WUORINEN AB; Iso
Roobertinkatu 4-6 A, FIN-00120 Helsinki (FI).

(21) International Application Number:

PCT/FI2003/000946

(22) International Filing Date:

11 December 2003 (11.12.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

20022178

11 December 2002 (11.12.2002)

FI

(71) Applicant (for all designated States except US): OY JURI-
LAB LTD [FI/FI]; Mikrokatu 1, FIN-70210 Kuopio (FI).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SALONEN,
Jukka, T. [FI/FI]; Kipinäkatu 4, FIN-70620 Kuopio
(FI). PIRSKANEN, Mia [FI/FI]; Kuninkaankatu 11 A
2, FI-70100 Kuopio (FI). TUOMAINEN, Tomi-Pekka
[FI/FI]; Louhikonkatu 35, FIN-70800 Kuopio (FI).
YUNUS, Faisal [PK/PK]; 10-F/A Model Town, 54000
Lahore (PK).

(81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE
(utility model), DE, DK (utility model), DK, DM, DZ, EC,
EE (utility model), EE, EG, ES, FI (utility model), FI, GB,
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL,
PT (utility model), PT, RO, RU, SC, SD, SE, SG, SK (util-
ity model), SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG,
US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD AND KIT FOR DETECTING A RISK FOR DIABETES OR A METABOLIC SYNDROME

(57) Abstract: The invention provides a method and kit for detecting or diagnosing a risk of or predisposition to diabetes or a metabolic syndrome in a subject, the method comprising the steps of providing a biological sample of the subject to be tested and detecting the presence or absence of a variant genotype of the human α_{2B} adrenoceptor in the biological sample. The presence of the variant genotype indicates an increased risk of diabetes or a metabolic syndrome in said subject. The invention also relates to a method for the treatment of type 2 diabetes.

WO 2004/053158 A1

Method and kit for detecting a risk for diabetes or a metabolic syndrome.

FIELD OF THE INVENTION

- 5 The invention provides a method and kit for detecting or diagnosing a risk of or predisposition to diabetes or a metabolic syndrome in a subject, the method comprising the steps of providing a biological sample of the subject to be tested and detecting the presence or absence of a variant genotype of the human α_{2B} -adrenoceptor in the biological sample. The invention also relates to a method for the treatment of diabetes.

10 BACKGROUND OF THE INVENTION

There are two sub-types of α -adrenoceptors: the α_1 -adrenoceptors (α_1 -ARs), on the effector organs (postsynaptic) and the α_2 -adrenoceptors (α_2 -ARs) on the nerve endings (presynaptic). The α_2 -adrenoceptors mediate many of the physiological effects of epinephrine and norepinephrine including a reduction of release of norepinephrine.

- 15 Three genetic subtypes of α_2 -ARs are known in humans, denoted as α_{2A} -, α_{2B} - and α_{2C} -AR that are located on chromosomes 10, 2 and 4, respectively (Calzada and Artinano 2001). The tissue distributions and physiological and pharmacological functions of the receptor subtypes have been reviewed elsewhere (Docherty 1998, Calzada and Artinano 2001). Based on recent studies with gene-targeted mice, α_{2A} -ARs mediate most of the
20 pharmacological actions ascribed to currently available α_2 -AR agonists, including inhibition of neurotransmitter release, central hypotensive and bradycardic effects, sedation and anaesthesia, and analgesia (MacMillan et al 1998).

- The pancreatic islets are richly innervated by parasympathetic, sympathetic and sensory nerves. Several different neurotransmitters are stored within the terminals of these
25 nerves, the classical neurotransmitters, acetylcholine and norepinephrine, and neuropeptides, which exert local effects. Stimulation of the autonomic nerves affects islet hormone secretion. Thus, insulin secretion is stimulated by parasympathetic nerves or their neurotransmitters and inhibited by sympathetic nerves or their neurotransmitters. The islet autonomic nerves seem to be of physiological importance

allowing oscillations of islet hormone secretion and are also involved in the islet adaptation in disease states (Ahrén 2000).

Sympathetic nervous system (SNS) over-activity plays a key role in the development of the metabolic syndrome, which is characterised by the combination of high blood pressure, increased glucose production, decreased glucose utilisation, increase in triglycerides and VLDL, decrease in HDL and insulin resistance (Borchard 2001).

Insulin resistance is defined as an impaired biologic response to exogenous or endogenous insulin and is a state in which the ability of insulin to suppress hepatic glucose output and stimulate the uptake and utilisation of glucose by muscle and adipose tissues is impaired (American Diabetes Association 1998).

The effects of the SNS on glucose and lipid metabolism are mediated by circulating catecholamines (epinephrine and norepinephrine) and the direct sympathetic innervation of the liver, adipose tissues and skeletal muscle. It has been generally recognised that increased sympathetic neural activity (SNA) produces catabolic effects on glucose and lipid metabolism (Nonogaki 2000) and inhibits insulin release from pancreatic beta-cells and glucose uptake into skeletal muscle cells, which are a major site of insulin-mediated glucose uptake (Borchard 2001) and a primary site of insulin resistance as measured by euglycaemic glucose clamp technique (Moan et al. 1996). Increased SNA causes vasoconstriction of the skeletal muscle arteries leading to impaired glucose tolerance and to insulin resistance.

In isolated adipocytes beta-adrenergic stimulation induces a rapid down-regulation of insulin receptors together with a decrease in insulin-mediated glucose transport. Insulin resistance leads to a breakdown of stored triglycerides in the adipose tissue and an increase in plasma free fatty acids. As a consequence, hepatic synthesis of triglycerides from free fatty acids and conversion of triglycerides to VLDL-cholesterol is enhanced. Catecholamines may further increase lipolysis in adipocytes which results in an elevated release of free fatty acids into the blood stream. Free fatty acids decrease glucose-stimulated insulin release from the pancreas, which further enhances glucose intolerance. Furthermore, catecholamines may inhibit lipoprotein lipase and thus increase VLDL, which is linked to a decrease in HDL (Borchard 2001). The end result

is a steady state of hyperinsulinemia. Support for this concept is based on these observations: a) a decreased skeletal muscle capillary density has been found in insulin-resistant states of hypertension, obesity, and type 2 diabetes; b) antihypertensive drugs that cause vasoconstriction worsen insulin resistance and those that cause vasodilation
5 improve insulin sensitivity; and c) exercise training improves insulin sensitivity and increases skeletal muscle capillary density (Julius et al. 1991).

The role of SNS in insulin resistance and diabetes is further strengthened by studies in which blockade of parasympathetic activity has been shown to increase the prevalence and severity of diabetes in mice suggesting role of unopposed sympathetic activity
10 (Kvist-Reimer et al. 2002) and sympathectomy abrogated the development of both hyperinsulinemia and hypertension in fructose hypertensive rats - a widely used model to study the inter-relationship between hyperinsulinemia, insulin resistance and high blood pressure (Verma et al. 1999).

In addition to the findings that increased SNA promotes insulin resistance and type 2
15 diabetes there is also experimental evidence that hyperinsulinemia may itself increase sympathetic neural outflow by activating anteroventral portion of the third ventricle in the central nervous system, a region implicated in arterial pressure regulation and sympathetic neural control. In rats and humans acute insulin elicits increased lumbar SNA along with elevated plasma norepinephrine and studies have firmly established in
20 both humans and experimental animals that insulin administration produces marked increases in SNA directed largely to skeletal muscle tissues (Muntzel 1999). Insulin is also involved in the initiation of appropriate changes in sympathetic outflow within the central nervous system in response to diet. During fasting state, the small decline in the plasma glucose concentration and the larger decrease in the plasma insulin
25 concentration result in diminished insulin-mediated uptake and metabolism of glucose to carbon dioxide by insulin-sensitive cells in the ventromedial hypothalamus. The decreased glucose metabolism stimulates the activity of an inhibitory pathway between the hypothalamus and the brain stem, suppressing tonically active sympathetic centres in the brain stem and decreasing central sympathetic activity. Conversely, in insulin-
30 resistant states, the increase in plasma glucose and the greater increase in plasma insulin (hyperinsulinemia) stimulate insulin-mediated uptake and metabolism of glucose by the hypothalamic cells.

The increase in glucose metabolism diminishes the activity of the inhibitory pathway, disinhibiting tonically active brain-stem centres and increasing central sympathetic activity leading to hypertension by stimulating the heart, the vasculature, and the kidneys (Reaven et al. 1996).

- 5 Thus it can be concluded that increased SNA and hyperinsulinemia by a mutually dependent process results in insulin resistance and increased risk of type 2 diabetes and carries a high mortality and risk of cardiovascular diseases, kidney failure and infectious complications as measured by resistance and pulsatility indices (Takahashi et al. 1998; Valensi et al. 1997; Weinrauch et al. 1995; Weston et al. 1999) and warrants
10 an early detection and treatment.

- Both β -adrenoceptor antagonists (β -blockers) and diuretics have been observed to influence glucose balance unfavourably, elevating blood glucose and serum insulin levels. Drug-induced hyperglycemia is a growing concern. Several antihypertensive drugs have an adverse effect on glucose tolerance that may partially or completely
15 negate the beneficial effects of reducing blood pressure as it relates to the incidence of coronary heart disease and its complications. Diuretics and beta-blockers have the greatest adverse effect on glucose intolerance (Houston 1986) and patients receiving beta-blocker treatment or diuretics may be at increased risk for developing hyperglycemia and subsequent diabetes mellitus (Luna et al. 2001; Bengtsson 1984;
20 Gress et al. 2000; Bengtsson et al. 1984).

- Metabolic studies of beta-blockers and diuretics aroused concern about the diabetogenic potential of these drugs. Subsequently, the results of prospective cohort studies and clinical trials suggested a causal link between the use of beta-blockers or diuretics and
25 the subsequent development of type 2 diabetes (Gress et al. 2000; Bengtsson et al. 1984; Lithell 1998; Papaccio et al. 1987). The risk of diabetes was 28 percent greater among those who took beta-blockers than among those who took no medication, without regard to the presence or absence of hypertension, sociodemographic characteristics, health-related behaviour, family history with respect to diabetes, and a
30 variety of coexisting conditions (Gress et al. 2000). Likewise, some observational studies have identified higher estimates of relative risk: for instance, in one study,

subjects who took beta-blockers had up to 6.1 times the risk of diabetes of those who did not (Samuelsson et al. 1994).

Thiazide diuretic-induced hyperglycemia occurs primarily through the reduction in total body potassium and the subsequent decreased insulin secretion (Luna et al. 2001; Greenberg 2000; Perez-Stable et al. 1983) by the beta cells, and reductions in extracellular fluid volume and cardiac output. This is compounded by increases in catecholamines from sympathetic nerve activity, which decrease peripheral glucose utilisation (Wilcox 1999). Potential mechanisms by which beta-blockers may contribute to the development of diabetes include weight gain, attenuation of the beta-receptor-mediated release of insulin from pancreatic beta cells, and decreased blood flow through the microcirculation in skeletal-muscle tissue, leading to decreased insulin sensitivity (Sowers et al. 2000).

SUMMARY OF THE INVENTION

The object of this invention is to provide a screening method to assess if an individual is at risk to develop diabetes or a metabolic syndrome based on the genotype of α_{2B} -adrenoceptor (ADRA2B) gene. The invention also provides a method for the treatment of diabetes in a human or animal subject. A further object of the invention is a method to determine whether a subject will benefit of different antihypertensive treatments with regard to their effects on the glucose balance and metabolism. Another object of the invention is a method for the selection of subjects for clinical trials testing antidiabetogenic drugs and compounds with effects on the insulin sensitivity.

The present invention concerns a method for detecting a risk of diabetes, especially type 2 diabetes, or a metabolic syndrome in a subject by determining the pattern of alleles encoding a variant α_{2B} -adrenoceptor i.e. to determine if said subject's genotype of the human α_{2B} -adrenoceptor is variant type, comprising the steps of

- a) providing a biological sample taken from the subject to be tested,

- b) detecting the presence or absence of variant genotype of the human α_{2B} -adrenoceptor gene in the biological sample, the presence of variant genotype indicating an increased risk of diabetes in said subject.

5 According to the invention, the method allows the determination whether said subject is of said variant genotype or not, a presence of said variant genotype in the biological sample, such as a blood sample or a buccal swab sample, thus indicating an increased risk of the subject to develop diabetes, and/or indicating the subject is in need for treatment, such as α_{2B} -selective or α_{2B} -nonselective antagonist therapy.

10 Preferably, the present invention concerns a method for detecting a risk of diabetes or a metabolic syndrome in a subject by determining the pattern of alleles encoding a variant α_2 -adrenoceptor gene i.e. to determine if said subject's genotype of the α_2 -adrenoceptor gene is of the deletion/deletion (D/D) type, comprising the steps of

- 15 a) providing a biological sample taken from the subject to be tested,
b) detecting the presence of α_2 -adrenoceptor deletion/deletion (D/D) type in the biological sample, the presence of D/D genotype indicating an increased risk of diabetes or a metabolic syndrome in said subject or the subject's need for α_{2B} -selective or α_{2B} -nonselective antagonist therapy for
20 diabetes.

A further object of the invention is a method for treating diabetes in a diabetic or glucose intolerant subject by reducing the sympathetic tone or activity, lowering blood or tissue norepinephrine or epinephrine concentrations and/or antagonizing α_{2B} -
25 adrenoceptors of the human or animal subject.

The present invention is also directed to a kit for detecting a risk of diabetes, especially type 2 diabetes, or a metabolic syndrome in a subject, comprising means for determining the pattern of alleles encoding a variant α_{2B} -adrenoceptor in a biological
30 sample.

In the spirit of the invention, it is also conceivable that any variation in the α_{2B} -adrenoceptor gene that alters the structure or function of the mature α_{2B} -adrenoceptor protein can be used in predicting the risk of diabetes or a metabolic syndrome, choosing a preferable or avoidable antihypertensive medication, or choosing a preferable or avoidable antidiabetic medication, or that said variant gene or materials connected thereto can be utilised as a part of a kit constructed for said uses.

DETAILED DESCRIPTION OF THE INVENTION

10 The present invention relates to a DNA molecule encoding a variant human α_{2B} -adrenoceptor, said variant α_{2B} -adrenoceptor protein and a method to assess the risk of individuals to develop diabetes or a metabolic syndrome in mammals as well as a method for the targeting treatment for diabetes and selecting subjects for clinical trials.

The word treating shall also be understood to include preventing.

15 The concept "a deletion of at least 1 glutamate from a glutamic acid repeat element of 12 glutamates" refers to any deletion of 1 to 12 glutamates, amino acids 298–309 (SEQ ID NO: 4), in an acidic stretch of 18 amino acids 294–311 located in the 3rd intracellular loop of the receptor polypeptide irrespective of the specific location of the deletion in said repeat element, or how many glutamates from said repeat element of 12
20 glutamates are deleted.

The concept "deletion/deletion (D/D) genotype of the human α_{2B} -adrenoceptor", in short "D/D genotype", refers to a genotype of an individual having both α_{2B} -adrenoceptor alleles code for a variant α_{2B} -adrenoceptor with a deletion of at least 1 glutamate from a glutamic acid repeat element of 12 glutamates, amino acids 298–309, in an acidic stretch of 18 amino acids 294–311 (SEQ ID NO: 4), located in the 3rd
25 intracellular loop of the receptor polypeptide. Correspondingly "deletion/insertion (D/I) genotype" refers to a genotype having one of the gene alleles code for an α_{2B} -adrenoceptor with a said deletion and the other without a said deletion, i.e. with a respective insertion, and thus the "insertion/insertion (I/I) genotype" refers to a

genotype having both alleles code for an α_{2B} -adrenoceptor without said deletion or deletions.

The term "metabolic syndrome" is defined in the Experimental Section.

A common variant form (SEQ ID NO: 1) of the human α_{2B} -AR gene (SEQ ID NO: 3) has been identified (Heinonen et al. 1999). This variant gene encodes a receptor protein (SEQ ID NO: 2) with a deletion of 3 glutamates, amino acids 307-309, from a glutamic acid (Glu) repeat element of 12 glutamates, amino acids 298-309, in an acidic stretch of 18 amino acids 294-311 (SEQ ID NO: 4), located in the 3rd intracellular loop of the receptor polypeptide. This variant gene (SEQ ID NO: 1) was associated with decreased basal metabolic rate (BMR) in a group of obese Finnish subjects (Heinonen et al. 1999). Of the 166 obese subjects, 47 (28 %) were homozygous for the long 12 glutamate repeat element (Glu¹²/Glu¹²), whereas 90 (54 %) were heterozygous (Glu¹²/Glu⁹) and 29 (17 %) were homozygous for the short form (Glu⁹/Glu⁹).

The results to be presented below show that in a population sample of Finnish middle-aged men subjects homozygous for the short form (Glu⁹/Glu⁹) described above, thus representing a deletion/deletion (D/D) genotype of the α_{2B} -adrenoceptor, have a significantly elevated risk for diabetes. Based on these results and previous publications referred to above it can be postulated that this D/D genotype is related to an impaired capacity to downregulate α_{2B} -adrenoceptor function during sustained receptor activation. Since altered α_{2B} -adrenoceptor function seems to be of relevance in the pathogenesis of diabetes, it could also be of relevance in subjects with the insertion/deletion (I/D) (heterozygous Glu¹²/Glu⁹) and insertion/insertion (I/I) (homozygous Glu¹²/Glu¹²) genotypes when other risk factors for diabetes are present. Further, since this specific deletion of 3 glutamates from said glutamic acid repeat element of 12 glutamates, amino acids 298-309, in said acidic stretch of 18 amino acids 294-311, located in the 3rd intracellular loop of the receptor polypeptide seems to be of relevance in diabetes then also other deletions, i.e. deletions of at least 1 glutamate, from said glutamic acid repeat element of 12 glutamates, amino acids 298-309, could be of relevance in the pathogenesis of diabetes, because the 3rd intracellular loop of the receptor polypeptide said repeat element is located in seems to have an essential role in

the down-regulation of the α_{2B} -adrenoceptor. Thus individuals with functional mutations in the α_{2B} -adrenoceptor gene have chronically up-regulated α_{2B} -adrenoceptors, leading to the vasoconstriction of peripheral arteries and reduced muscle and pancreatic blood flow.

- 5 α_{2B} -adrenoceptors mediate contraction of arteries, and genetic polymorphism present in the α_{2B} -adrenoceptor gene renders some subjects more susceptible to α_{2B} -adrenoceptor mediated vasoconstriction of the peripheral blood flow regulating arteries and arterioles and associated clinical disorders such as diabetes or a metabolic syndrome. These subjects will especially benefit from treatment with an α_{2B} -adrenoceptor antagonist, and
- 10 will be at increased risk for adverse effects if subtype-nonselective α_2 -agonists are administered to them. Therefore, a gene test recognizing subjects with a deletion variant of the α_{2B} -adrenoceptor gene will be useful in diagnostics and patient selection for specific therapeutic procedures and clinical drug testing trials. A gene test recognizing the D/D genotype of the α_{2B} -adrenoceptor is useful in assessing an individual's risk to
- 15 develop diabetes related to the D/D genotype. The test can be used to set a specific subdiagnosis of diabetes, based on its genetic etiology.

Furthermore, a gene test recognizing the D/D genotype of the α_{2B} -adrenoceptor is useful in selecting drug therapy for patients with diabetes. Such drugs are e.g. a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic

20 receptors of the subjects either directly or through central nervous system effects, for example pindolol, propranolol, sotalol, timolol, acebutolol, atenol, betaxolol, bisoprol, esmolol, metoprolol, seliprol, carvedilol, labetalol, clonidine, moxonidine, prazosin, or indapamid, including α -adrenoceptor antagonists (α_{2B} -selective or nonselective).

- 25 For instance, as angiotensin II causes an increase of noradrenaline sensitivity, and this effect is at least in part mediated by α -adrenoceptors (Datte et al. 2000), the blood pressure lowering effect of drugs acting through angiotensin II inhibition, such as the angiotensin (AT) receptor blockers, is conceivably enhanced in persons with the D/D genotype of the α_{2B} -adrenoceptor. Such drugs are for
- 30 example captopril, cinapril, enalapril, imidapril, lisinopril, moexipril, perindopril,

ramipril, trandolapril, candesartan, eprosartan, irbesartan, losartan, valsartan or telmisartan.

5 A gene test recognizing the D/D genotype of the α_{2B} -adrenoceptor is useful in selecting drug therapy for patients who might be at increased risk for adverse effects of α_2 -adrenergic agonists; either it will be possible to avoid the use of α_2 -agonists in such patients, or it will be possible to include a specific α_{2B} -antagonist in their therapeutic regimen.

10 For instance, a nucleic acid sample from a subject can be used for determining if said subject is a carrier of a variant gene. The determination can be carried out either as a DNA analysis according to well known methods, which include direct DNA sequencing of the normal and variant gene, allele specific amplification using the polymerase chain reaction (PCR) enabling detection of either normal or variant sequence, or by indirect detection of the normal or variant gene by various molecular biology methods including e.g. PCR-single stranded conformation polymorphism (SSCP) or conformational analysis (SSCA) method or denaturing gradient gel electrophoresis (DGGE). Determination of the normal or variant gene can also be done by using a restriction fragment length polymorphism (RFLP) method, which is particularly suitable for genotyping large numbers of samples. Similarly, a test based on gene chip or array technology can be easily developed in analogy with many currently existing such tests for single-nucleotide polymorphisms.

25 The determination can also be carried out at the level of RNA by analyzing RNA expressed at tissue level using various methods. Allele specific probes can be designed for hybridization. Hybridization can be done e.g. using Northern blot, RNase protection assay or in situ hybridization methods. RNA derived from the normal or variant gene can also be analyzed by converting tissue RNA first to cDNA and thereafter amplifying cDNA by an allele specific PCR method.

The presence of variant α_{2B} -adrenoceptor polypeptides indicating the presence of variant gene can be detected by various methods such as hybridization (Western blot), or other antibody based protein assays.

A kit for detecting a risk of diabetes or a metabolic syndrome preferably contains the various components needed for carrying out the diagnostic method according to the present invention. These components are preferably packaged in separate containers and/or vials. The kit may also include instructions for carrying out the method. Thus, 5 for example, some or all of the various reagents and other ingredients needed for carrying out the determination, such as buffers, primers, enzymes, control samples or standards etc. can be packaged separately but provided for use in the same box.

Instructions for carrying out the method can be included inside the box, as a separate insert, or as a label on the box and/or on the separate vials. The kit may also contain the 10 necessary computer software needed to interpret the results obtained with the kit, or for utilizing the results from a gene chip used in the method. Preferably, the kit contains a capturing nucleic acid or an antibody specific to a variant α_{2B} -adrenoceptor nucleic acid or polypeptide, respectively.

15 The publications and other materials used herein to illuminate the background of the invention, and in particular, to provide additional details with respect to its practice, are incorporated herein by reference.

The invention will be described in more detail in the experimental section.

EXPERIMENTAL SECTION

Determination of genomic alleles encoding the α_{2B} -adrenoceptor

DNA fragment analysis of ADRA2B insertion/deletion polymorphism

5 ADRA2B insertion/deletion mutation

The nucleotide sequence of the primer pair for the amplification of the human ADRA2B gene (alpha2B-adrenergic receptor gene) insertion/deletion polymorphism (SEQ ID NO:3) (SEQ ID NO:1) was as follows 5'- GGG TGT TTG TGG GGC ATC TC -3' (SEQ ID NO:5) and 5'- TGG CAC TGC CTG GGG TTC A -3' (SEQ ID
10 NO:6). A fluorescent label has been added to the 5' end of one of the above mentioned PCR primers. Thus, the pcr fragment is detectable in the capillary electrophoresis conducted with ABI Prism 3100 Genetic Analyzer.

The genomic DNA region of the mutation in question can be amplified with PCR with
15 PTC-220 DNA Dyad PCR machine (MJ Research). The PCR reaction was conducted in a 20 μ l volume: the reaction mixture contained 60 ng human genomic DNA (extracted from peripheral blood), 1X PCR Buffer (QIAGEN), 100 μ l of each of the nucleotides (dATP, dCTP, dGTP, dTTP), 0.5 μ M of each of the primers and 1 unit of the DNA polymerase (QIAGEN, Hot Start Taq DNA polymerase). The PCR conditions need to
20 be determined experimentally, and the following standard protocol can be used as a start: first the reaction was hold 7 minutes at 94°C, then the following three steps were repeated for 35 cycles: 45 secondes at 94°C, 45 secondes at 54°C, 1 minute at 72°C, after which the reaction was kept at 72°C for an additional 5 minutes and finally hold at 4°C.

25

The insertion/deletion polymorphism of ADRA2B gene concerns an insertion or an deletion of three glutamic acids in the region of 12 Glu aminoacids in the codons 298-309 (SEQ ID NO:3). Thus depending on the genotype, there is either 9 Glu (deletion) (SEQ ID NO:2) or 12 Glu (insertion) (SEQ ID NO:4) at the ADRA2B protein.
30 Depending on whether the amplified allele had an insertion or a deletion in the studied

locus, the size of the PCR product was 91 bp (insertion allele) or 82 bp (deletion allele). Thus, for homozygotes (insertion/insertion or deletion/deletion) only one size of a fragment was detected either 91 bp or 82 bp, respectively. For heterozygotes both of the above mentioned fragments were detected.

5

Capillary electrophoresis with ABI Prism 3100 Genetic Analyzer

A sample of the ADRA2B insertion/deletion PCR product, 9.00 µl of Hi-Di formamide (Applied Biosystems) and 0.25 µl GeneScan-120 LIZ size standard (Applied Biosystems) were combined in a 96-well 3100 optical microamp plate (Applied Biosystems). The reactions were denatured by placing them at 95°C for 5 minutes and then loaded onto an ABI Prism 3100 Genetic Analyzer (Applied Biosystems). Electrophoresis data was processed and the genotypes were visualized by using the GeneScan Analysis version 3.7 (Applied Biosystems).

Study Population

15 The above referred study population of 1426 Finnish middle-aged men subjects including 304 subjects with a specific deletion/deletion (D/D) genotype of the α_{2B} -adrenoceptor is described in more detail in the following:

Knowing the possible involvement of the investigated acidic region in the desensitization mechanism of the receptor we hypothesized that the observed
20 insertion/deletion allelic variation could be associated with insulin resistance, as indicated by serum insulin concentration, the metabolic syndrome and type 2 diabetes. To test this hypothesis, we carried out a population study in middle-aged Finnish men. The study was carried out as part of the Kuopio Ischemic Heart Disease Risk Factor Study (KIHD), which is an ongoing population-based study designed to investigate risk
25 factors for cardiovascular diseases, type 2 diabetes and related outcomes in men from eastern Finland (Salonen 1988). This area is known for its homogenous population (Sajantila et al. 1996) and high coronary morbidity and mortality rates (Keys 1980).

Body-mass index was computed as the ratio of weight (kg) to the square of height (m). Waist circumference was taken as the average of 2 measurements taken after inspiration
30 and after expiration (mean difference between the two measurements \cong 1.5cm) at the

midpoint between the lowest rib and the iliac crest. Waist-hip ratio was defined as the ratio of waist girth to the circumference of the hips measured at the trochanter major.

Subjects were asked to fast for 12 h before blood sampling. They were also asked to refrain from smoking for 12 h and from consuming alcohol for 3 days before blood
5 draws. Blood glucose was measured at baseline and 11-year follow-up using a glucose dehydrogenase method after precipitation of proteins by trichloroacetic acid. The serum samples for insulin determination were stored at -80°C. Serum insulin was determined with a Novo Biolabs radioimmunoassay kit (Novo Nordisk, Bagsvaerd, Denmark).

Diabetes was defined as fasting blood glucose of 6.7 mmol/L or more or diagnosed
10 diabetes with either dietary, oral or insulin treatment. The metabolic syndrome for men according to the WHO definition was modified for epidemiological studies as proposed by the EGIR (Balkau et al. 1999) and defined as: hyperinsulinemia (fasting insulin levels in the top 25% of the non-diabetic population), impaired fasting glycemia or diabetes and the presence of at least two of the following: abdominal obesity,
15 dyslipidemia (triglycerides ≥ 1.70 or HDL < 0.9 mmol/l), or hypertension (blood pressure $\geq 140/90$ mm Hg or blood pressure medication) (Alberti et al 1998). Insulin resistance was approximated as hyperinsulinemia based on fasting insulin concentrations in the upmost fourth (Balkau et al. 1999). Insulin resistance was also estimated as the bottom fourth of insulin sensitivity as measured by a validated index
20 (QUICKI) based on fasting insulin and glucose concentrations ($[\log(\text{insulin}) + \log(\text{glucose})]^{-1}$) (Katz et al 2000). Hypertension was defined according to the EGIR recommendations at a lower level than the original WHO definition for consistence with current WHO-ISH and Sixth Joint National Committee recommendations (Balkau et al. 1999, The sixth report 1997, Guidelines Subcommittee 1999). Microalbuminuria
25 was not included in the definition (Balkau et al. 1999).

Abdominal obesity was defined according to two definitions: 1) according to the original WHO definition - waist-hip ratio > 0.90 or body-mass index ≥ 30 kg/m² (Alberti et al 1998) and 2) modified according to the EGIR recommendation - waist circumference ≥ 94 cm (Balkau et al. 1999).

The metabolic syndrome as defined by the NCEP comprises three or more of the following indications: fasting plasma glucose levels ≥ 6.1 mmol/l (blood glucose levels ≥ 5.6 mmol/l), serum triglycerides ≥ 1.7 mmol/l, serum HDL < 1.0 mmol/l, blood pressure $\geq 135/85$ mmHg, waist girth > 102 cm (Executive Summary 2001). Use of waist girth > 94 cm was suggested for men genetically susceptible to insulin resistance (Executive Summary 2001).

Of all the 1426 subjects, 304 (21%) had the homozygous deletion (D/D) genotype. Of 809 subjects who had no diabetes at the baseline examination and for whom 11-year follow-up information was available, 168 were D/D homozygotes. Of the subjects with D/D genotype, 19 (11.3%) developed diabetes during 11 years of follow-up, whereas of the other subjects, 39 (6.1%) developed diabetes. The odds ratio was 2.0 (95% confidence interval 1.1 to 3.5, $p=0.028$ in 2-sided Fisher's exact test and $p=0.021$ in a logistic model). Other strongest predictors of diabetes during the follow-up were body-mass index (kg/m²) and the waist-to-hip circumference ratio. In a multivariate logistic model including these covariates, the D/D genotype was associated with a 1.9-fold (95% confidence interval 1.1 to 3.5, $p=0.033$) probability (incidence) of diabetes.

The mean fasting baseline serum insulin was 11.1 (mU/L) in 304 D/D homozygotes and 10.5 mU/L in other 1122 genotyped men ($p=0.045$ for difference in Mann-Whitney U test). The D/D genotype of α_{2B} -adrenoceptor gene was also associated with the prevalence and incidence of metabolic syndrome.

The effects of α -adrenoceptor antagonists, β -adrenoceptor antagonists (beta-blockers) and diuretics on serum insulin and lipids and diabetes was analyzed separately in 278 subjects with an antihypertensive medication, of whom 66 were α_{2B} -AR deletion/deletion homozygotes and 212 had other genotypes. Among the non-homozygotes (the wild I allele carriers), the use of α -adrenoceptor antagonists was associated with a lowering of both fasting (6.9 vs. 11.3 mU/L, $p=0.020$) and 2h post glucose load serum insulin concentrations (34.0 vs. 72.3 mU/L, $p<0.001$), while among the D/D homozygotes, α -adrenoceptor antagonists had no effect on either fasting or post-load serum insulin concentration. Among the α_{2B} -AR deletion homozygotes, the use of β -adrenoceptor antagonist therapy was associated with a lowering of serum total

(5.1 vs 5.8 mmol/L, $p=0.010$) and LDL cholesterol (3.6 vs. 4.1 mmol/L, $p=0.053$) concentrations, while among other genotypes, both serum cholesterol and LDL cholesterol tended to be higher among β -blocker users than among non-users. Among the deletion homozygotes, the use of diuretics was associated elevations of serum total
5 (6.0 vs 5.5 mmol/L) and LDL cholesterol (4.4 vs 3.8 mmol/L) and lowering of triglycerides (1.5 vs 2.1 mmol/L, $p=0.005$), whereas among subjects with other genotypes, serum total and LDL cholesterol tended to be lower and triglycerides higher among the diuretic users.

The prevalence of diabetes (fasting blood glucose of 6.0 mmol/L or more or diagnosed
10 diabetes with either dietary, oral or insulin treatment) at the KIHHD 11-year follow-up was in the α_{2B} -AR deletion carriers higher among α -adrenoceptor antagonist users than non-users (25.0% vs 13.2%), while in the non-carriers, the prevalence of diabetes was lower among the users than the non-users (0% vs 12.3%).

Taken together, the known biological properties of the α_{2B} -AR, the homogeneity of the
15 Finnish population, the study design, the relatively large representative study population and the association of diabetes with one trait suggest that the D/D receptor allele is a causal genetic risk factor for diabetes and modifies the effects of α - and β -adrenoceptor modulating drugs and diuretics on glucose and lipid metabolism. The α_{2B} -AR deletion allele homozygosity or carrier status appears to attenuate or abolish the
20 beneficial effect of α -adrenoceptor antagonists on glucose metabolism, to induce cholesterol and LDL cholesterol elevation by diuretics and reverse or attenuate the cholesterol and LDL cholesterol elevating effect of β -blocking agents.

It will be appreciated that the methods of the present invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will
25 be apparent for the specialist in the field that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are illustrative and should not be construed as restrictive.

References

- Ahren B. Autonomic regulation of islet hormone secretion--implications for health and disease. *Diabetologia*. 2000; 43: 393-410.
- 5 Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539-53.
- American Diabetes Association. Consensus Development Conference on Insulin Resistance. *Diabetes Care*. 1998; 21: 310-314.
- 10 Balkau B, Charles MA. Comment on the provisional report from the WHO consultation. European Group for the Study of Insulin Resistance (EGIR). *Diabet Med* 1999;16:442-3.
- 15 Bengtsson C. Metabolic side effects of diuretics and beta-adrenoceptor blockers. *Acta Pharmacol Toxicol*. 1984; 54 Suppl 1: 67-70.
- Bengtsson C, Blohme G, Lapidus L, Lindquist O, Lundgren H, Nystrom E, Petersen K, Sigurdsson JA. Do antihypertensive drugs precipitate diabetes? *Br Med J (Clin Res Ed)*. 1984; 289: 1495-1497.
- 20 Borchard U. The role of the Sympathetic Nervous System in Cardiovascular Disease. *J Clin Basic Cardiol*. 2001; 4: 175-177.
- 25 Calzada BC, Artinano AL. Alpha-adrenoceptor subtypes. *Pharmacol Res* 2001; 44: 195-208.
- Docherty JR. Subtypes of functional α_1 - and α_2 -receptors. *Eur J Pharmacol* 1998; 361: 1-15.
- 30 1-15.

Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). JAMA 2001;285:2486-97.

- 5 Greenberg A. Diuretic complications. Am J Med Sci. 2000; 319: 10-24. Perez-Stable E, Caralis PV. Thiazide-induced disturbances in carbohydrate, lipid, and potassium metabolism. Am Heart J. 1983; 106: 245-251.

- 10 Gress TW, Nieto FJ, Shahar E, Wofford MR, Brancati FL. Hypertension and antihypertensive therapy as risk factors for type 2 diabetes mellitus. Atherosclerosis Risk in Communities Study. N Engl J Med. 2000; 342: 905-912.

- 15 Heinonen P, Koulu M, Pesonen U, Karvonen M, Rissanen A, Laakso M, Valve R, Uusitupa M, Scheinin M. Identification of a three amino acid deletion in the alpha-2B-adrenergic receptor which is associated with reduced basal metabolic rate in obese subjects. J Clin Endocrinol Metab 1999;84:2429-2433.

- 20 Houston MC. Adverse effects of antihypertensive drug therapy on glucose intolerance. Cardiol Clin. 1986; 4: 117-135.

Jousilahti P, Vartiainen E, Tuomilehto J, Pekkanen J, Puska P. Role of known risk factors in explaining the difference in the risk of coronary heart disease between eastern and southwestern Finland. Ann Med 1998; 30: 481-487.

- 25 Julius S, Gudbrandsson T, Jamerson K, Tariq Shahab S, Andersson O. The hemodynamic link between insulin resistance and hypertension. J Hypertens. 1991; 9: 983-986.

Kaplan GA, Salonen JT. Socioeconomic conditions in childhood and ischaemic heart disease during middle age. BMJ 1990; 301: 1121-1123.

Katz A, Nambi SS, Mather K, et al. Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 2000;85:2402-10.

- 5 Keys A. Seven Countries: A Multivariate Analysis of Death and Coronary Heart Disease. Cambridge, Mass, Harvard University Press, 1980.

Kvist-Reimer M, Sundler F, Ahren B. Effects of chemical sympathectomy by means of 6-hydroxydopamine on insulin secretion and islet morphology in alloxan-diabetic mice. *Cell Tissue Res.* 2002; 307: 203-209.

10

Laukkanen JA, Kurl S, Lakka TA, et al. Exercise-induced silent myocardial ischemia and coronary morbidity and mortality in middle-aged men. *J Am Coll Cardiol* 2001;38(1):72-79.

Lithell HO. Insulin resistance and diabetes in the context of treatment of hypertension.

- 15 Blood Press. 1998; Suppl 3: 28-31.

Luna B, Feinglos MN. Drug-induced hyperglycemia. *JAMA.* 2001; 286: 1945-1948.

- 20 MacMillan LB, Lakhani P, Lovinger D, Limbird LE. Alpha 2-adrenergic receptor subtypes: subtle mutation of the alpha 2A-adrenergic receptor in vivo by gene targeting strategies reveals the role of this subtype in multiple physiological settings. *Recent Prog Horm Res.* 1998; 53: 25-42.

- 25 Moan A, Eide IK, Kjeldsen SE. Metabolic and adrenergic characteristics of young men with insulin resistance. *Blood Press Suppl.* 1996; 1: 30-37.

Muntzel MS. Insulin-mediated sympathoexcitation in obesity and type 2 diabetes. *Nephrol Dial Transplant.* 1999; 14: 2282-2285.

Nonogaki K. New insights into sympathetic regulation of glucose and fat metabolism. *Diabetologia*. 2000; 43: 533-549.

5 Papaccio G, Esposito V. Hyperglycemic effects of hydrochlorothiazide and propranolol. A biochemical and ultrastructural study. *Acta Diabetol Lat*. 1987; 24: 325-330.

10 Reaven GM, Lithell H, Landsberg L. Hypertension and associated metabolic abnormalities--the role of insulin resistance and the sympathoadrenal system. *N Engl J Med*. 1996; 334: 374-381.

Sajantila A, Salem AH, Savolainen P, Bauer K, Gierig C, Paabo S: Paternal and maternal DNA lineages reveal a bottleneck in the founding of the Finnish population. *Proc Natl Acad Sci U.S.A.* 1996;93:12035-12039.

15 Salonen JT. Is there a continuing need for longitudinal epidemiologic research? The Kuopio Ischaemic Heart Disease Risk Factor Study. *Ann Clin Res* 1988;20:46-50.

20 Samuelsson O, Hedner T, Berglund G, Persson B, Andersson OK, Wilhelmsen L. Diabetes mellitus in treated hypertension: incidence, predictive factors and the impact of non-selective beta-blockers and thiazide diuretics during 15 years treatment of middle-aged hypertensive men in the Primary Prevention Trial Goteborg, Sweden. *J Hum Hypertens* 1994; 8: 257-263.

25 Sowers JR, Bakris GL. Antihypertensive therapy and the risk of type 2 diabetes mellitus. *N Engl J Med*. 2000; 342: 969-970.

Takahashi T, Nishizawa Y, Emoto M, Kawagishi T, Matsumoto N, Ishimura E, Inaba M, Okuno Y, Shimada H, Morii H. Sympathetic function test of vasoconstrictor changes in foot arteries in diabetic patients. *Diabetes Care*. 1998; 21: 1495-1501.

30 The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med* 1997;157:2413-46.

Valensi P, Huard JP, Giroux C, Attali JR. Factors involved in cardiac autonomic neuropathy in diabetic patients. *J Diabetes Complications*. 1997; 11:180-187.

- 5 Verma S, Bhanot S, McNeill JH. Sympathectomy prevents fructose-induced hyperinsulinemia and hypertension. *Eur J Pharmacol*. 1999; 373: R1-4.

Weinrauch LA, D'Elia JA, Gleason RE, Keough J, Mann D, Kennedy FP. Autonomic function in type I diabetes mellitus complicated by nephropathy. A cross-sectional
10 analysis in the presymptomatic phase. *Am J Hypertens*. 1995; 8: 782-789.

Weston PJ, Gill GV. Is undetected autonomic dysfunction responsible for sudden death in Type 1 diabetes mellitus? The 'dead in bed' syndrome revisited. *Diabet Med*. 1999; 16: 626-631.

15

Wilcox CS. Metabolic and adverse effects of diuretics. *Semin Nephrol*. 1999 ; 19: 557-568.

1999 World Health Organization-International Society of Hypertension Guidelines for
20 the Management of Hypertension. Guidelines Subcommittee. *J Hypertens* 1999;17:151-83.

CLAIMS

1. A method for detecting a risk of diabetes or a metabolic syndrome in a subject by determining the pattern of alleles encoding a variant α_{2B} -adrenoceptor protein, comprising the steps of
 - a) providing a biological sample taken from the subject to be tested,
 - b) detecting the presence or absence of a variant genotype of the human α_{2B} -adrenoceptor in the biological sample, the presence of the variant genotype indicating an increased risk of diabetes or a metabolic syndrome in said subject.
2. The method according to claim 1, where said variant genotype of the human α_{2B} -adrenoceptor gene is any nucleotide variation in the genomic DNA that affects the structure or function of the mature α_{2B} -adrenoceptor protein.
3. The method according to claim 2, wherein said variant genotype of the human α_{2B} -adrenoceptor is a homozygote deletion form of mutation (D/D).
4. The method according to claim 1, wherein the detection step is a DNA-assay.
5. The method according to claim 1, wherein the detection step is carried out using a gene or DNA chip, microarray, strip, panel or similar combination of more than one genes, mutations or RNA expressions to be assayed.
6. The method according to claim 1, wherein the allelic pattern is determined using polymerase chain reaction.
7. The method according to claim 1, wherein the biological sample is any sample that comprises genetic material, preferably a blood sample or buccal swab sample.

8. The method according to claim 1, wherein the detection step is based on a capturing probe, a single strand of cDNA, comprising a nucleotide sequence encoding a variant α_{2B} -adrenoceptor protein or a fragment thereof.
9. The method according to claim 1, wherein the detection step is an analysis
5 of the α_{2B} -adrenoceptor polypeptide or a part thereof.
10. The method according to claim 1, wherein said method is used for determining whether a subject will benefit from a treatment with a drug affecting the norepinephrine sensitivity or sympathetic activity of the subject.
11. The method according to claim 1, wherein said method is used for
10 determining whether a subject will benefit from a treatment with an α - or α_{2B} -adrenoceptor antagonist.
12. The method according to claim 11, wherein said method is used for determining whether the treatment with an α - or α_{2B} -adrenoceptor antagonist will lower or elevate serum insulin concentration.
- 15 13. The method according to claim 1, wherein said method is used for determining whether a subject will be at increased risk of hyperglycemia, hyperinsulinemia, a metabolic syndrome and diabetes if a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptors or a diuretic is administered to the subject.
- 20 14. The method according to claim 1, wherein said method is used for determining whether a subject will response to a treatment with diuretics or a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptors.
- 25 15. The method according to claim 1, wherein said method is used for determining whether the treatment with diuretics and a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptor will lower or elevate serum and plasma cholesterol, LDL cholesterol and triglyceride concentrations.

16. The method according to any one of claims 13 - 15, wherein said drug is pindolol, propranolol, sotalol, timolol, acebutolol, atenol, betaxolol, bisoprol, esmolol, metoprolol, seliprol, carvedilol, labetalol, clonidine, moxonidine, prazosin, or indapamid.

5 17. The method according to claim 1 further comprising a step of selecting a subject of the D/D genotype for clinical drug trials testing antidiabetic and insulin sensitivity improving effects of compounds.

18. The method according to claim 17, wherein the compound to be tested is a drug affecting the norepinephrine sensitivity or sympathetic activity of the
10 subject.

19. The method according to claim 17, wherein the compound to be tested is a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenergic receptors of the subject either directly or through central nervous system.

20. The method according to claim 1, wherein said diabetes is type 2 diabetes.

15 21. The method according to claim 1, wherein said metabolic syndrome is defined as hyperinsulinemia (fasting insulin levels in the top 25% of the non-diabetic population), impaired fasting glycemia or diabetes and the presence of at least two of the following: abdominal obesity, dyslipidemia (triglycerides ≥ 1.70 or HDL < 0.9 mmol/l), or hypertension (blood pressure $\geq 140/90$ mm Hg or blood
20 pressure medication).

22. A method for treating a human or animal subject suffering from type 2 diabetes or for treating vascular complications of diabetes, said method comprising a step of reducing the sympathetic tone, lowering blood or tissue norepinephrine or epinephrine concentrations and/or antagonizing α - or α_{2B} -adrenoceptors of the human or animal
25 subject.

23. The method according to claim 21, wherein said method comprise a step of administering to a subject a compound reducing the sympathetic tone, lowering blood

or tissue norepinephrine or epinephrine concentrations and/or antagonizing α_{2B} -adrenoceptors of the human subject or animal.

24. The method according to claim 23, wherein said compound is a subtype of
5 a selective or nonselective α_{2B} -adrenoceptor antagonist.

25. The method according to claim 23 or 24, wherein said method is a gene therapy or gene transfer method.

26. The method according to claim 25, wherein the gene to be transferred is a non-variant α_{2B} -adrenoceptor gene or a fragment or derivative thereof.

10 27. A kit for detecting a risk of diabetes or a metabolic syndrome in a subject, comprising means for determining the pattern of alleles encoding a variant α_{2B} -adrenoceptor in a biological sample taken from said subject, and optionally computer software to interpret the results of the determination.

28. The kit according to claim 27 for determining the presence or absence of a
15 variant genotype of the human α_{2B} -adrenoceptor in said biological sample.

29. The kit according to claim 27 or 28 comprising a capturing nucleic acid probe or an antibody.

SEQUENCE LISTING

<110> Oy Jurilab Ltd

<120> Method and kit for detecting a risk of diabetes

<130> 40548

<150> FI 20022178

<151> 2002-12-11

<160> 6

<170> PatentIn version 3.1

<210> 1

<211> 1344

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1344)

<223> ADRA2B variant type sequence

```

<400> 1
atg gac cac cag gac ccc tac tcc gtg cag gcc aca gcg gcc ata gcg      48
Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala
1          5          10          15

gcg gcc atc acc ttc ctc att ctc ttt acc atc ttc ggc aac gct ctg      96
Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu
          20          25          30

gtc atc ctg gct gtg ttg acc agc cgc tgc ctg cgc gcc cct cag aac      144
Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn
          35          40          45

ctg ttc ctg gtg tgc ctg gcc gcc gcc gac atc ctg gtg gcc acg ctc      192
Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu
          50          55          60

```

atc atc cct ttc tcg ctg gcc aac gag ctg ctg ggc tac tgg tac ttc Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe 65 70 75 80	240
cgg cgc acg tgg tgc gag gtg tac ctg gcg ctc gac gtg ctc ttc tgc Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys 85 90 95	288
acc tcg tcc atc gtg cac ctg tgc gcc atc agc ctg gac cgc tac tgg Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp 100 105 110	336
gcc gtg agc cgc gcg ctg gag tac aac tcc aag cgc acc ccg cgc cgc Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg 115 120 125	384
atc aag tgc atc atc ctc act gtg tgg ctc atc gcc gcc gtc atc tcg Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser 130 135 140	432
ctg ccg ccc ctc atc tac aag ggc gac cag ggc ccc cag ccg cgc ggg Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly 145 150 155 160	480
cgc ccc cag tgc aag ctc aac cag gag gcc tgg tac atc ctg gcc tcc Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser 165 170 175	528
agc atc gga tct ttc ttt gct cct tgc ctc atc atg atc ctt gtc tac Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr 180 185 190	576
ctg cgc atc tac ctg atc gcc aaa cgc agc aac cgc aga ggt ccc agg Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg 195 200 205	624
gcc aag ggg ggg cct ggg cag ggt gag tcc aag cag ccc cga ccc gac Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp 210 215 220	672
cat ggt ggg gct ttg gcc tca gcc aaa ctg cca gcc ctg gcc tct gtg His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val 225 230 235 240	720
gct tct gcc aga gag gtc aac gga cac tcg aag tcc act ggg gag aag Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys 245 250 255	768
gag gag ggg gag acc cct gaa gat act ggg acc cgg gcc ttg cca ccc Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro 260 265 270	816
agt tgg gct gcc ctt ccc aac tca ggc cag ggc cag aag gag ggt gtt Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val 275 280 285	864
tgt ggg gca tct cca gag gat gaa gct gaa gag gag gaa gag gag gag Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu 290 295 300	912

3

gag gag tgt gaa ccc cag gca gtg cca gtg tct ccg gcc tca gct tgc 960
 Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala Ser Ala Cys
 305 310 315 320

agc ccc ccg ctg cag cag cca cag ggc tcc cgg gtg ctg gcc acc cta 1008
 Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu Ala Thr Leu
 325 330 335

cgt ggc cag gtg ctc ctg ggc agg ggc gtg ggt gct ata ggt ggg cag 1056
 Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile Gly Gly Gln
 340 345 350

tgg tgg cgt cga agg gcg cac gtg acc cgg gag aag cgc ttc acc ttc 1104
 Trp Trp Arg Arg Ala His Val Thr Arg Glu Lys Arg Phe Thr Phe
 355 360 365

gtg ctg gct gtg gtc att ggc gtt ttt gtg ctc tgc tgg ttc ccc ttc 1152
 Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp Phe Pro Phe
 370 375 380

ttc ttc agc tac agc ctg ggc gcc atc tgc ccg aag cac tgc aag gtg 1200
 Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His Cys Lys Val
 385 390 395 400

ccc cat ggc ctc ttc cag ttc ttc ttc tgg atc ggc tac tgc aac agc 1248
 Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr Cys Asn Ser
 405 410 415

tca ctg aac cct gtt atc tac acc atc ttc aac cag gac ttc cgc cgt 1296
 Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp Phe Arg Arg
 420 425 430

gcc ttc cgg agg atc ctg tgc cgc ccg tgg acc cag acg gcc tgg tga 1344
 Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr Ala Trp
 435 440 445

<210> 2

<211> 447

<212> PRT

<213> Homo sapiens

<400> 2

Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala
 1 5 10 15

Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu
 20 25 30

Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn
 35 40 45

Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu
 50 55 60

Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe
 65 70 75 80

Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys
 85 90 95

Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp
 100 105 110

Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg
 115 120 125

Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser
 130 135 140

Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly
 145 150 155 160

Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser
 165 170 175

Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr
 180 185 190

Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg
 195 200 205

Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp
 210 215 220

His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val
 225 230 235 240

Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys
 245 250 255

Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro
 260 265 270

Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val
 275 280 285

Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu

5

290

295

300

Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala Ser Ala Cys
 305 310 315 320

Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu Ala Thr Leu
 325 330 335

Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile Gly Gly Gln
 340 345 350

Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg Phe Thr Phe
 355 360 365

Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp Phe Pro Phe
 370 375 380

Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His Cys Lys Val
 385 390 395 400

Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr Cys Asn Ser
 405 410 415

Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp Phe Arg Arg
 420 425 430

Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr Ala Trp
 435 440 445

<210> 3

<211> 1353

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (1353)

<223> ADRA2B wild type sequence

<400> 3

atg gac cac cag gac ccc tac tcc gtg cag gcc aca gcg gcc ata gcg

Met	Asp	His	Gln	Asp	Pro	Tyr	Ser	Val	Gln	Ala	Thr	Ala	Ala	Ile	Ala	
1				5					10					15		
gcg	gcc	atc	acc	ttc	ctc	att	ctc	ttt	acc	atc	ttc	ggc	aac	gct	ctg	96
Ala	Ala	Ile	Thr	Phe	Leu	Ile	Leu	Phe	Thr	Ile	Phe	Gly	Asn	Ala	Leu	
			20					25					30			
gtc	atc	ctg	gct	gtg	ttg	acc	agc	cgc	tcg	ctg	cgc	gcc	cct	cag	aac	144
Val	Ile		Leu	Ala	Val	Leu	Thr	Ser	Arg	Ser	Leu	Arg	Ala	Pro	Gln	Asn
		35						40				45				
ctg	ttc	ctg	gtg	tcg	ctg	gcc	gcc	gcc	gac	atc	ctg	gtg	gcc	acg	ctc	192
Leu	Phe	Leu	Val	Ser	Leu	Ala	Ala	Ala	Asp	Ile	Leu	Val	Ala	Thr	Leu	
	50					55					60					
atc	atc	cct	ttc	tcg	ctg	gcc	aac	gag	ctg	ctg	ggc	tac	tgg	tac	ttc	240
Ile	Ile	Pro	Phe	Ser	Leu	Ala	Asn	Glu	Leu	Leu	Gly	Tyr	Trp	Tyr	Phe	
65					70					75					80	
cgg	cgc	acg	tgg	tgc	gag	gtg	tac	ctg	gcg	ctc	gac	gtg	ctc	ttc	tgc	288
Arg	Arg	Thr	Trp	Cys	Glu	Val	Tyr	Leu	Ala	Leu	Asp	Val	Leu	Phe	Cys	
				85					90					95		
acc	tcg	tcc	atc	gtg	cac	ctg	tgc	gcc	atc	agc	ctg	gac	cgc	tac	tgg	336
Thr	Ser	Ser	Ile	Val	His	Leu	Cys	Ala	Ile	Ser	Leu	Asp	Arg	Tyr	Trp	
			100					105					110			
gcc	gtg	agc	cgc	gcg	ctg	gag	tac	aac	tcc	aag	cgc	acc	ccg	cgc	cgc	384
Ala	Val	Ser	Arg	Ala	Leu	Glu	Tyr	Asn	Ser	Lys	Arg	Thr	Pro	Arg	Arg	
		115					120					125				
atc	aag	tgc	atc	atc	ctc	act	gtg	tgg	ctc	atc	gcc	gcc	gtc	atc	tcg	432
Ile	Lys	Cys	Ile	Ile	Leu	Thr	Val	Trp	Leu	Ile	Ala	Ala	Val	Ile	Ser	
	130					135					140					
ctg	ccg	ccc	ctc	atc	tac	aag	ggc	gac	cag	ggc	ccc	cag	ccg	cgc	ggg	480
Leu	Pro	Pro	Leu	Ile	Tyr	Lys	Gly	Asp	Gln	Gly	Pro	Gln	Pro	Arg	Gly	
145					150					155					160	
cgc	ccc	cag	tgc	aag	ctc	aac	cag	gag	gcc	tgg	tac	atc	ctg	gcc	tcc	528
Arg	Pro	Gln	Cys	Lys	Leu	Asn	Gln	Glu	Ala	Trp	Tyr	Ile	Leu	Ala	Ser	
				165					170					175		
agc	atc	gga	tct	ttc	ttt	gct	cct	tgc	ctc	atc	atg	atc	ctt	gtc	tac	576
Ser	Ile	Gly	Ser	Phe	Phe	Ala	Pro	Cys	Leu	Ile	Met	Ile	Leu	Val	Tyr	
			180					185					190			
ctg	cgc	atc	tac	ctg	atc	gcc	aaa	cgc	agc	aac	cgc	aga	ggg	ccc	agg	624
Leu	Arg	Ile	Tyr	Leu	Ile	Ala	Lys	Arg	Ser	Asn	Arg	Arg	Gly	Pro	Arg	
		195					200					205				
gcc	aag	ggg	ggg	cct	ggg	cag	ggg	gag	tcc	aag	cag	ccc	cga	ccc		

7

245	250	255	
gag gag ggg gag acc cct gaa gat act ggg acc cgg gcc ttg cca ccc Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro			816
260	265	270	
agt tgg gct gcc ctt ccc aac tca ggc cag ggc cag aag gag ggt gtt Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val			864
275	280	285	
tgt ggg gca tct cca gag gat gaa gct gaa gag gag gaa gag gag gag Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu			912
290	295	300	
gag gag gag gaa gag tgt gaa ccc cag gca gtg cca gtg tct ccg gcc Glu Glu Glu Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala			960
305	310	315	320
tca gct tgc agc ccc ccg ctg cag cag cca cag ggc tcc cgg gtg ctg Ser Ala Cys Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu			1008
325	330	335	
gcc acc cta cgt ggc cag gtg ctc ctg ggc agg ggc gtg ggt gct ata Ala Thr Leu Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile			1056
340	345	350	
ggg ggg cag tgg tgg cgt cga agg gcg cac gtg acc cgg gag aag cgc Gly Gly Gln Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg			1104
355	360	365	
ttc acc ttc gtg ctg gct gtg gtc att ggc gtt ttt gtg ctc tgc tgg Phe Thr Phe Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp			1152
370	375	380	
ttc ccc ttc ttc ttc agc tac agc ctg ggc gcc atc tgc ccg aag cac Phe Pro Phe Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His			1200
385	390	395	400
tgc aag gtg ccc cat ggc ctc ttc cag ttc ttc ttc tgg atc ggc tac Cys Lys Val Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr			1248
405	410	415	
tgc aac agc tca ctg aac cct gtt atc tac acc atc ttc aac cag gac Cys Asn Ser Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp			1296
420	425	430	
ttc cgc cgt gcc ttc cgg agg atc ctg tgc cgc ccg tgg acc cag acg Phe Arg Arg Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr			1344
435	440	445	
gcc tgg tga Ala Trp			1353
450			

<210> 4

<211> 450

<212> PRT

<213> Homo sapiens

<400> 4

Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala
 1 5 10 15

Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu
 20 25 30

Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn
 35 40 45

Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu
 50 55 60

Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe
 65 70 75 80

Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys
 85 90 95

Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp
 100 105 110

Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg
 115 120 125

Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser
 130 135 140

Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly
 145 150 155 160

Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser
 165 170 175

Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr
 180 185 190

Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg
 195 200 205

Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp
 210 215 220

His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val
225 230 235 240

Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys
245 250 255

Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro
260 265 270

Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val
275 280 285

Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu
290 295 300

Glu Glu Glu Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala
305 310 315 320

Ser Ala Cys Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu
325 330 335

Ala Thr Leu Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile
340 345 350

Gly Gly Gln Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg
355 360 365

Phe Thr Phe Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp
370 375 380

Phe Pro Phe Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His
385 390 395 400

Cys Lys Val Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr
405 410 415

Cys Asn Ser Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp
420 425 430

Phe Arg Arg Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr
435 440 445

Ala Trp
450

<210> 5

<211> 20

<212> DNA ..

<213> Artificial

<220>

<221> misc_feature

<222> (1)..(20)

<223> ADRA2B pcr primer f

<400> 5

gggtgtttgt ggggcatctc

20

<210> 6

<211> 19

<212> DNA

<213> Artificial

<220>

<221> misc_feature

<222> (1)..(19)

<223> ADRA2B pcr primer r

<400> 6

tggcactgcc tggggttca

19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 2003/000946

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12Q 1/68 // C07K 14/72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, BIOSIS, MEDLINE, EMBASE, CHEM.ABS.DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02066617 A2 (OY JURILAB LTD), 29 August 2002 (29.08.2002), page 9, line 16 - line 25, claim 22	27-29
A	--	1-21
A	The Journal of Clinical Endocrinology & Metabolism, Volume 84, 1999, No. 7, Paula Heinson et al, "Identification of a Three-Amino Acid Deletion in the alpha 2B - Adrenergic Receptor That Is Associated with Reduced Basal Metabolic Rate in Obese Subjects", pages 2429-2433, abstract; page 2432	1-21,27-29
	--	

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

11 February 2004

Date of mailing of the international search report

19-02-2004

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

TERESE PERSSON/BS

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.
PCT/FI 2003/000946

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item I.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:

a. type of material

- ☒ a sequence listing
- ☐ table(s) related to the sequence listing

b. format of material

- ☒ in written format
- ☒ in computer readable form

c. time of filing/furnishing

- ☒ contained in the international application as filed
- ☒ filed together with the international application in computer readable form
- ☐ furnished subsequently to this Authority for the purposes of search

2. ☒ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 2003/000946

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	International Journal of Obesity, Volume 25, 2001, K. Sivenius et al, "Effect of a three-amino acid deletion in the alpha2B-adrenergic receptor gene on long-term body weight change in Finnish non-diabetic and type 2 diabetic subjects", pages 1609-1614, abstract --	1-21,27-29
A	US 2002058618 A1 (WURSTER ET AL), 16 May 2002 (16.05.2002), abstract; page 1, column 1, paragraph 4 - column 2, paragraph 3 --	1-21,27-29
A	US 6150389 A (MUNK ET AL), 21 November 2000 (21.11.2000), abstract, table 1 --	1-21,27-29
A	Annu. Rev. Pharmacol. Toxicol., Volume 32, 1993, Robert R. Ruffolo et al, "Pharmacologic and therapeutic applications of alpha2-adrenoceptor subtypes", pages 243-79 --	1-21,27-29
P,A	Clinical Autonomic Research, Volume 13, 2003, Gerasimos P. Sykiotis et al, "The alpha2B adrenergic receptor deletion/insertion polymorphism in morbid obesity", pages 203-207, abstract --	1-21,27-29
P,X	International Journal of Obesity, Volume 27, 2003, Suppl. 1, Siitonen N. et al, "The Association of Polymorphism in the alpha2B- Adrenoceptor Gene with Type 2 Diabetes and Obesity", Column T2:P2-017, abstract -- -----	1-9,20

INTERNATIONAL SEARCH REPORT

Information on patent family members

24/12/2003

International application No.

PCT/FI 2003/000946

WO	02066617	A2	29/08/2002	EP	1368454 A	10/12/2003
				FI	20010323 A,V	21/08/2002
				US	2003003470 A	02/01/2003

US	2002058618	A1	16/05/2002	US	6521632 B	18/02/2003
				AU	3551001 A	20/08/2001
				BR	0108221 A	05/03/2003
				CA	2399421 A	16/08/2001
				CZ	20022884 A	12/02/2003
				EE	200200435 A	15/12/2003
				EP	1253926 A	06/11/2002
				HU	0300032 A	28/05/2003
				IL	151017 D	00/00/0000
				JP	2003522148 T	22/07/2003
				NO	20023773 A	09/08/2002
				SK	11472002 A	04/02/2003
				WO	0158454 A	16/08/2001

US	6150389	A	21/11/2000	US	6319935 B	20/11/2001
				US	6569884 B	27/05/2003
				US	2002049241 A	25/04/2002
				US	2003187047 A	02/10/2003
				US	5731337 A	24/03/1998
				AU	687584 B	26/02/1998
				AU	3007695 A	09/02/1996
				CA	2194702 A	25/01/1996
				DE	69524223 D,T	22/08/2002
				EP	0770068 A,B	02/05/1997
				ES	2168373 T	16/06/2002
				JP	10502643 T	10/03/1998
				WO	9601813 A	25/01/1996

INTERNATIONAL SEARCH REPORT

International application No.
PCT/FI 2003/000946

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **22-26**
because they relate to subject matter not required to be searched by this Authority, namely:
see extra sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/FI 2003/000946

Box II.1

Claims 22-26 relate to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practiced on the human or animal body (PCT Rule 39.1(iv)). No search has been carried out for these claims.